



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

155. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, O.

A bought a horse, which he sold to B at a loss of $m=6\%$; B sold the horse to C at a loss of $n=5\%$; and C sold the horse to D at a gain of $p=12\frac{1}{2}\%$. How much did A lose, if C gained $\$G=\26.79 ?

ALGEBRA.

156. Proposed by B. F. FINKEL, A. M., M. Sc., Professor of Mathematics and Physics in Drury College, Springfield, Mo.

$(z+x)a - (z-x)b = 2yz \dots (1)$; $(x+y)b - (x-y)c = 2xz \dots (2)$; $(y+z)c - (y-z)a = 2xy \dots (3)$. Find the values of x , y , and z , by the method of linear simultaneous equations.

NOTE. This problem was somewhat abbreviated in the last issue. The problem occurs in Fisher & Schwatt's *Elements of Algebra*, page 224, under Linear Simultaneous Equations.

157. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics in The Temple College, Philadelphia, Pa.

Solve the equations,

$$\frac{x}{a+\lambda} + \frac{y}{b+\lambda} + \frac{z}{c+\lambda} + \frac{u}{d+\lambda} = 1, \quad \frac{x}{a+\mu} + \frac{y}{b+\mu} + \frac{z}{c+\mu} + \frac{u}{d+\mu} = 1,$$

$$\frac{x}{a+\nu} + \frac{y}{b+\nu} + \frac{z}{c+\nu} + \frac{u}{d+\nu} = 1, \quad \frac{x}{a+\rho} + \frac{y}{b+\rho} + \frac{z}{c+\rho} + \frac{u}{d+\rho} = 1.$$

158. Proposed by R. D. BOHANNON, Ph. D., Professor of Mathematics, Ohio State University, Columbus, O.

$$\text{If } \frac{x}{a+a} + \frac{y}{b+a} + \frac{z}{c+a} = \frac{x}{a+\beta} + \frac{y}{b+\beta} + \frac{z}{c+\beta} = \frac{x}{a+\gamma} + \frac{y}{b+\gamma} + \frac{z}{c+\gamma} = 1,$$

$$\text{show, } \frac{x}{(a+\beta)^2} + \frac{y}{(b+\beta)^2} + \frac{z}{(c+\beta)^2} = \frac{(\gamma-\beta)(a-\beta)}{(a+\beta)(b+\beta)(c+\beta)}.$$

GEOMETRY.

182. Proposed by MARCUS BAKER, U. S. Geological Survey, Washington, D. C.

Show how to cut from a given cube, edge s , the maximum tetrahedron.

183. Proposed by S. F. NORRIS, Professor of Mathematics and Astronomy, Baltimore City College, Baltimore, Md.

"Two quadrilaterals having three sides of the one equal to the three corresponding sides of the other, each to each, and the two corresponding angles adjacent to the unknown sides equal, each to each, are equal figures." [Olney's *Geometry*, page 129.]